

- i. Office of Statewide Health Planning and Development, Patient Discharge Data
- ii. Healthy Stores for a Healthy Community Assessment, San Mateo County raw data, 2016.
- iii. Substance Abuse and Mental Health Services Administration (SAMHSA), HHS, Results from the 2015 National Survey on Drug Use and Health, NSDUH: Detailed Tables, 2016. <http://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2015/NSDUH-DetTabs-2015/NSDUHDetTabs-2015.pdf>
- iv. Ambrose, B.K., et al., Flavored Tobacco Product Use Among US Youth Aged 12-17 Years, 2013-2014. *JAMA*, 2015: p. 1-3
- v. FDA, Preliminary Scientific Evaluation of the Possible Public Health Effects of Menthol versus Nonmenthol Cigarettes, 2013
- vi. U.S. Department of Health and Human Services. 2012. Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General. Atlanta: U.S. National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, p. 539, www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf.
- vii. King BA, Tynan MA, Dube SR, et al. 2013. "Flavored-Little-Cigar and Flavored-Cigarette Use Among U.S. Middle and High School Students." *Journal of Adolescent Health*. [Article in press], www.jahonline.org/article/S1054-139X%2813%2900415-1/abstract.
- viii. Singh T, Arrazola RA, Corey CG, et al. Tobacco Use Among Middle and High School Students — United States, 2011– 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65(14):361-367. doi:10.15585/mmwr.mm6514a1.
- ix. Substance Abuse and Mental Health Services Administration. *The NSDUH Report: Use of Menthol Cigarettes*. 2009. <https://perma.cc/4UG9-HUQ2>.
- x. Fallin A, Goodin AJ, King BA. Menthol cigarette smoking among lesbian, gay, bisexual, and transgender adults. *Am J Prev Med*. 2015;48(1):93-97. doi:10.1016/j.amepre.2014.07.044.
- xi. Cohn AM, Johnson AL, Hair E, Rath JM, Villanti AC. Menthol tobacco use is correlated with mental health symptoms in a national sample of young adults: implications for future health risks and policy recommendations. *Tob Induc Dis*. 2016;14(1):1. doi:10.1186/s12971-015-0066-3.
- xii. Cantrell J, Kreslake JM, Ganz O, et al. Marketing little cigars and cigarillos: Advertising, price, and associations with neighborhood demographics. *Am J*

Public Health. 2013;103(10):1902-1909. doi:10.2105/AJPH.2013.301362.

- xiii. Villanti AC, Mowery PD, Delnevo CD, Niaura RS, Abrams DB, Giovino DB. Changes in the prevalence and correlates of menthol cigarette use in the USA, 2004-2014. *Tob Control*. 2016;25:ii14-ii20. doi: 10.1136/tobaccocontrol-2016-053329.
- xiv. Levy DT, Pearson JL, Villanti AC, et al. Modeling the future effects of a menthol ban on smoking prevalence and smoking-attributable deaths in the United States. *Am J Public Health*. 2011;101(7):1236-1240. doi:10.2105/AJPH.2011.300179.
- xv. Katz MH. 2008. "Banning Tobacco Sales in Pharmacies: The Right Prescription." *Journal of the American Medical Association*, 300(12):1451-1453.
- xvi. Tobacco Education and Research Oversight Committee for California. 2006. *Confronting a Relentless Adversary: A Plan for Success Toward a Tobacco-Free California 2006-2008*, p. 18, www.cdph.ca.gov/programs/tobacco/Documents/TEROC/2006-2008%20Master%20Plan.pdf.